

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 January 2003 (23.01.2003)

PCT

(10) International Publication Number  
**WO 03/006475 A2**

(51) International Patent Classification<sup>7</sup>: **C07H 19/00**

(21) International Application Number: PCT/DK02/00488

(22) International Filing Date: 12 July 2002 (12.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
PA 2001 01095 12 July 2001 (12.07.2001) DK

(71) Applicant (for all designated States except US):  
**CUREON A/S** [DK/DK]; Fruebjergvej 3, DK-2100  
Copenhagen Ø (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KOCH, Troels**


[DK/DK]; Funkiavej 47, DK-2300 Copenhagen S (DK).  
**ROSENBOHM, Christoph** [DK/DK]; Frederikssundsvej  
55, 4. th., DK-2400 Copenhagen NV (DK). **PEDERSEN,  
Daniel, Sejer** [DK/DK]; Knabrostræde 22, 3., DK-1210  
Copenhagen K (DK).

(74) Agent: **PLOUGMANN & VINGTOFT A/S**; Sundkrogs-  
gade 9, P.O. Box 831, DK-2100 Copenhagen Ø (DK).


(81) Designated States (*national*): AE, AG, AL, AM, AT (util-  
ity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (util-  
ity model), DE, DK (utility model), DK, DM, DZ, EC, EE  
(utility model), EE, ES, FI (utility model), FI, GB, GD, GE,  
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,  
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,  
MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD,  
SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR,  
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

[Continued on next page]


(54) Title: METHOD FOR PREPARATION OF LNA PHOSPHORAMIDITES




1



2



3



4

(57) Abstract: The present invention relates to large scale preparation of LNA phosphoramidites using a 2-cyanoethyl-N,N,N',N'-tetra-substituted phosphoramidite and a nucleophilic activator, e.g. 2-cyanoethyl-N,N,N',N'-tetraiso-propylphosphoramidite and 4,5-dicyanoimidazole. The method is faster and more cost efficient than previously known methods.



WO 03/006475 A2



**(84) Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## METHOD FOR PREPARATION OF LNA PHOSPHORAMIDITES

### FIELD OF THE INVENTION

The present invention relates to a high yielding, fast method for large-scale LNA phosphoramidites synthesis.

### 5 BACKGROUND OF THE INVENTION

Locked Nucleic Acid (LNA) monomers and oligonucleotides were invented in 1997 (WO9914226) and have showed promising results as antisense drug candidates. However, there is no production method described in the literature for large scale LNA-phosphoramidite synthesis providing optimal yields of the LNA amidites necessary for  
10 efficient preparation of LNA based antisense drugs.

Oligonucleotide synthesis is typically performed using the phosphoramidite method, invented by Caruthers, US4415732, in 1980 and improved a couple of years later by Köster US4725677. LNA oligomers are being synthesised according to the phosphoramidite  
15 method, but so far the large scale supply of LNA monomers has been a problem due to slow reactions, side product formation during the reaction and reagents unstable at room temperature have been employed.

The mechanism of phosphoramidite activation and coupling in oligonucleotide syntheses  
20 has been studied in detail. Traditionally tetrazole has been used (Dahl *et al.* 1987, *Nucleic Acid Research*, 15, 1729-1743; Beaucage & Iver, 1992, *Tetrahedron*, 48, 2223-2311). The proposed mechanism of tetrazole is two step, first tetrazole protonates trivalent phosphorous followed by displacement of the *N,N*-diisopropylamine by the tetrazolide. This latter intermediate is very reactive with hydroxynucleophiles such as 5'-OH on nucleic  
25 acids. Therefore, tetrazole acts both as acid and as nucleophilic agent.

Vargeese *et al.* (Vargeese, C.; Carter, J.; Krivjansky, S.; Settle, A.; Kropp, E.; Peterson, K.; Pieken, W. *Nucleic Acid Research*, 1998, 26, 1046-1050; WO9816540;) have described an activator for the coupling of phosphoramidites to the 5'-hydroxyl group during  
30 oligonucleotide synthesis. The activator is 4,5-dicyanoimidazole (DCI) and its effectiveness is thought to be based on its nucleophilicity. It was shown that DCI significantly increased the yield in phosphoramidite oligomerisation.

Vargeese *et al.* has described a method of small scale DNA phosphoramidite thymine  
35 monomer synthesis using DCI, but the method they used provided a moderate yield (75%) after several re-precipitations of the amidite. Kittaka *et al.* (Kittaka A., Kuze T.; Amano M.;

Tanaka H.; Miyasaka T.; Hirose K.; Yoshida T.; Sarai A.; Yasukawa T. and Ishii S.

*Nucleosides & Nucleotides* 18, 2769-2783, 1999) have also used DCI for amidite synthesis

but they reported an even lower yield of the product (65%). Kittaka *et al.* (Kittaka A.; Horii C.; Kuze T.; Asakura T.; Ito K.; Nakamura K.T.; Miyasaka T. and Inoue J., *Synthetic*

5 *letters*, S1, 869-872, 1999) have also made a derivatised uridine phosphoramidite by 3'-O phosphitylation using DCI (0.7 eq), but without describing the reaction parameters further. The yield was reported to be 92%. Generally, phosphitylation of uracil and thymine provide the highest yields because these nucleobases usually do not interfere with the phosphitylation reagents.

10

Accordingly, there is nothing in the literature describing that DCI has been used as an activator for LNA phosphoramidite synthesis.

The previously reported oxy- $\beta$ -D-ribo-LNA (Figure 2) method of synthesis used 2-

15 cyanoethyl-*N,N*-diisopropylchlorophosphoramidite and diisopropylethylamine (Koshkin, A. A.; Singh, S. K.; Nielsen, P.; Rajwanshi, V. K.; Kumar, R.; Meldgaard, M.; Olsen, C. E.; Wengel, J. *Tet.* 1998, 54, 3607-3630). The reported yields were low (LNA-T 70%, LNA-U

58%, LNA-G 64%, LNA-A 73% and no yield for LNA-C was reported). The previously reported method for synthesising the oxy- $\alpha$ -L-ribo thymine phosphoramidite (Figure 3)

20 also utilized 2-cyanoethyl-*N,N*-diisopropylchlorophosphoramidite and diisopropylethylamine and gave a yield of 60% (Hakansson, A.E.; Koshkin, A.A.; Sorensen M.D.; Wengel J. *J.Org.Chem.* 2000, 65, 5161-5166).

## BRIEF DESCRIPTION OF THE FIGURES

Figure 1: Phosphitylation of guanine, a) 2-Cyanoethyl-*N,N*-diisopropylchlorophosphor-

25 amidite, diisopropylethylamine, b) Phosphitylation of guanine 3'OH via the transfer/-rearrangement reaction (slow).

Figure 2: Oxy- $\beta$ -D-ribo-LNA phosphoramidites.

30 Figure 3: Amine- $\beta$ -D-ribo-LNA phosphoramidites and oxy- $\alpha$ -L-ribo-LNA thymine phosphoramidite.


## SUMMARY OF THE INVENTION

The present invention provides a novel high yielding and fast method for the large scale

35 synthesis of pure LNA phosphoramidites. In the production of LNA monomers as phosphoramidites, the phosphitylation step is crucial. If necessary also a facile and efficient purification is desired, due to the reactivity of the phosphoramidites, so as to

avoid their decomposition during purification. An absolute purity of the amidites is a prerequisite because the monomers are used in long sequential oligomerisations over some time where also impurities can cause the decomposition of the phosphoramidites.

- 5 The novel strategy is demonstrated by the synthesis of oxy- $\beta$ -ribo-LNA phosphoramidites where B is A<sup>Bz</sup>, MeC<sup>Bz</sup>, G<sup>Ibu</sup> and T and oxy  $\alpha$ -L-ribo-LNA-T (Figure 3).



- 10 The present invention provides a method for the synthesis of an LNA phosphoramidite, the method comprising phosphitylation of the 3'-OH groups of an LNA monomer with a 2-cyanoethyl-*N,N,N',N'*-tetra-substituted phosphoramidite in the presence of a nucleophilic activator.
- 15 The advantages of the present activation method is:
- a) Easy handling
  - b) All reagents are stabile at room temperature
  - c) Only 0.7 equivalents DCI are necessary
  - d) Easy workup
- 20 e) Fast and high yielding reaction at room temperature
- f) Cost efficient reaction
  - g) Chromatography not essential for obtaining amidites of high purity

The known phosphitylation side reaction of the guanine nucleobase followed by slow transfer/rearrangement to give the desired amidite (Nielsen, J.; Taagaard, M.; Marugg, J. E.; van Boom, J. H.; Dahl, O. *Nuc.Acid.Res.* 1986, 14, 7391-7403) (see Figure 1) was not observed using the conditions according to the present invention.

## DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "LNA monomer" refers to a ribonucleotide having a 2',4'-bridge (in particular a -O-CH<sub>2</sub>- (oxy-LNA), -S-CH<sub>2</sub>- (thio-LNA), -NR-CH<sub>2</sub>- (amino-LNA, R being

30 hydrogen, C<sub>1-6</sub>-alkyl, phenyl, benzyl, etc.) bridge) as described in the International Patent

Application WO9914226 and subsequent WO0056746, WO0056748, WO0066604 and WO0228875.

Particularly interesting examples of LNA monomers are those referred to as oxy- $\beta$ -D-ribo-LNA (cf. Figure 2), thio- $\beta$ -D-ribo-LNA, amino- $\beta$ -D-ribo-LNA, oxy- $\alpha$ -L-ribo-LNA (cf. Figure 3), thio- $\alpha$ -L-ribo-LNA or amino- $\alpha$ -L-ribo-LNA.

As used herein, the term "LNA phosphoramidite" refers to a nucleotide with a protected 5'-hydroxy group (e.g. DMT protected as illustrated in Figures 2 and 3), in which the 3'-hydroxy group is coupled to a trivalent phosphorous atom, which in turn is bonded to a suitable leaving group such as a *N,N*-dialkylamine, e.g. diisopropylamine and a protecting group such as the cyanoethyl (NCCH<sub>2</sub>CH<sub>2</sub>-) group. Any LNA phosphoramidite that can be used in solid or liquid phase oligonucleotide synthesis can be used in the present invention, including protected dimer and trimer LNA phosphoramidite.

As used herein, the term "2-cyanoethyl-*N,N,N',N'*-tetra-substituted phosphoramidite", also referred to as "PN<sub>2</sub>-reagent", refers in short to a 2-cyanoethyl-*N,N,N',N'*-tetra-substituted phosphoramidite, wherein the term "substituted" means a linear, cyclic or branched unsaturated hydrocarbon such as allyl or vinyl or saturated hydrocarbon or substituted hydrocarbon group or aromatic group or substituted aromatic group having 1 to 9 carbon atoms, such as methyl, ethyl, propyl, iso-propyl, pentyl, cyclopentyl, hexyl, cyclohexyl, preferred examples alkyl are methyl, ethyl, propyl, iso-propyl, butyl, tert-butyl, iso-butyl, pentyl, cyclopentyl, hexyl, cyclohexyl, in particular methyl, ethyl, propyl, iso-propyl, tert-butyl, iso-butyl, cyclohexyl, propylene, butylene, pentylene, hexylene, heptylene, where the substituents can be a heteroatom such as oxygen or nitrogen.

The LNA monomer as well as the phosphoramidite LNA monomer may carry any nucleobase in the 1' position.

In the present context, the term "nucleobase" covers naturally occurring nucleobases as well as non-naturally occurring nucleobases. It should be clear to the person skilled in the art that various nucleobases which previously have been considered "non-naturally occurring" have subsequently been found in nature. Thus, "nucleobase" includes not only the known purine and pyrimidine heterocycles, but also heterocyclic analogues and tautomers thereof. Illustrative examples of nucleobases are adenine, guanine, thymine, cytosine, uracil, purine, xanthine, diaminopurine, 8-oxo-*N*<sup>6</sup>-methyladenine, 7-deazaxanthine, 7-deazaguanine, *N*<sup>4</sup>,*N*<sup>4</sup>-ethanocytosine, *N*<sup>6</sup>,*N*<sup>6</sup>-ethano-2,6-diaminopurine, 5-methylcytosine, 5-(C<sup>3</sup>-C<sup>6</sup>)-alkynylcytosine, 5-fluorouracil, 5-bromouracil, pseudoisocytosine, 2-hydroxy-5-methyl-4-triazolopyridin, isocytosine, isoguanin, inosine, *N*<sup>6</sup>-

alylpurines, *N*<sup>6</sup>-acylpurines, *N*<sup>6</sup>-benzylpurine, *N*<sup>6</sup>-halopurine, *N*<sup>6</sup>-vinylpurine, *N*<sup>6</sup>-acetylenic  
 purine, *N*<sup>6</sup>-acyl purine, *N*<sup>6</sup>-hydroxyalkyl purine, *N*<sup>6</sup>-thioalkyl purine, *N*<sup>2</sup>-alkylpurines, *N*<sup>4</sup>-  
 alkylpyrimidines, *N*<sup>4</sup>-acylpyrimidines, *N*<sup>4</sup>-benzylpurine, *N*<sup>4</sup>-halopyrimidines, *N*<sup>4</sup>-  
 vinylpyrimidines, *N*<sup>4</sup>-acetylenic pyrimidines, *N*<sup>4</sup>-acyl pyrimidines, *N*<sup>4</sup>-hydroxyalkyl  
 5 pyrimidines, *N*<sup>6</sup>-thioalkyl pyrimidines, thymine, cytosine, 6-azapyrimidine, including 6-  
 azacytosine, 2- and/or 4-mercaptopyrimidine, uracil, *C*<sup>5</sup>-alkylpyrimidines, *C*<sup>5</sup>-  
 benzylpyrimidines, *C*<sup>5</sup>-halopyrimidines, *C*<sup>5</sup>-vinylpyrimidine, *C*<sup>5</sup>-acetylenic pyrimidine, *C*<sup>5</sup>-  
 acyl pyrimidine, *C*<sup>5</sup>-hydroxyalkyl purine, *C*<sup>5</sup>-amidopyrimidine, *C*<sup>5</sup>-cyanopyrimidine, *C*<sup>5</sup>-  
 nitropyrimidine, *C*<sup>5</sup>-aminopyrimidine, *N*<sup>2</sup>-alkylpurines, *N*<sup>2</sup>-alkyl-6-thiopurines, 5-  
 10 azacytidinyl, 5-azauracilyl, trazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, and  
 pyrazolopyrimidinyl. Functional oxygen and nitrogen groups on the base can be protected  
 as necessary or desired. Suitable protecting groups are well known to those skilled in the  
 art, and included trimethylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, and *t*-  
 butyldiphenylsilyl, trityl, alkyl groups, acyl groups such as acetyl and propionyl,  
 15 methanesulfonyl, and *p*-toluenesulfonyl, imines such as *N*-dimethylmethyletheneamine  
 and dibutylmethyletheneamine. Preferred bases include cytosine, methyl cytosine, uracil,  
 thymine, adenine and guanine.

LNA phosphoramidites according to the present invention are prepared in high yield with 2-  
 20 cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite (a PN<sub>2</sub>-reagent). This reagent has not  
 successfully been used to prepare LNA phosphoramidites before. Previously 2-cyanoethyl-  
*N,N*-diisopropylchlorophosphoramidite has been used for preparation of LNA amidites. This  
 has not afforded the high purity, high yields and low reaction times according to the  
 present invention.

25 The surprising finding in the present inventions is that 2-cyanoethyl-*N,N,N',N'*-tetraiso-  
 propylphosphoramidite can be activated by 4,5-dicyanoimidazole and that this complex  
 provides the accurate reactivity, during the conditions described, to secure efficient  
 phosphorylation of all 4 LNA DMT-protected nucleosides including the G-nucleoside. With  
 30 the previous known methods using 2-cyanoethyl-*N,N*-diisopropylchlorophosphoramidite the  
 phosphorylation reaction would proceed via oxygen on the protected guanine nucleobases  
 (Figure 1a) and thereafter be rearranged to the thermodynamically stable 3'-O-amidite  
 (Figure 1b). The disadvantages with the previous method were that the final product  
 contained side products and that it required significantly longer reaction time, (up to 24 h)  
 35 due to the slow rearrangement resulting in the thermodynamical product.

The traditional procedures for making LNA phosphoramidites provided yields ranging from  
 14–73%, whereas the present procedure provides yields of 95% or higher. This is also

remarkable for the reason that the corresponding reaction for DNA/RNA nucleotides only provides yields of 92%.

Another advantage of the present invention is that the reactions are finished in 0.5-4

5 hours, preferably 1.0-3.5 hours, and that chromatography of the LNA amidites is unnecessary due to the efficient reaction. Thus, the phosphitylation reaction is typically allowed to proceed for 0.1-12 hours, such as 0.2-8 hours, e.g. 0.5-4.0 hours, such as 1.0-3.5 hours.

10 An LNA phosphoramidite for direct use in oligonucleotide synthesis may be obtained directly or by simple precipitation of the product from the reaction mixture. Thus, in a particular aspect of the invention, the method comprises at the most one precipitation step.

15 Suitable nucleophilic activators for use in the present invention include, but are not limited to, 4,5-dicyanoimidazole (DCI), 4-alkylthioimidazole, 2-alkylthioimidazole, 2-nitroimidazole, 4-nitroimidazole, 4,5-dihaloimidazole, 4-haloimidazole, 2-haloimidazole and 5-alkoxytetrazole. DCI is the preferred nucleophilic activator. DCI has a  $pK_a$  of 5.2 and is easily dissolved in acetonitrile.

20

Suitable  $PN_2$ -reagents are for example, but not limited to, 2-cyanoethyl-*N,N,N',N'*-tetra-isopropylphosphoramidite and 2-cyanoethyl-*N,N,N',N'*-tetraethylphosphoramidite and the reagents described by Dahl, B.H., Nielsen J. and Dahl O. *Nucleic Acid Research*, 1987, 15, 1729-1743 all incorporated here by reference. It is presently believed that 2-cyanoethyl-

25 *N,N,N',N'*-tetraisopropylphosphoramidite and 2-cyanoethyl-*N,N,N',N'*-

tetraethylphosphoramidite are especially relevant, in particular 2-cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite.

A currently preferred combination is where the 2-cyanoethyl-*N,N,N',N'*-tetra-substituted

30 phosphoramidite is the 2-cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite and the nucleophilic activator is 4,5-dicyanoimidazole.

The molar ratio between the LNA monomer and the nucleophilic activator is typically in the range of 1:0.0001 to 1:10, preferably 1:0.0001 to 1:1, and more preferably 1:0.0001 to

35 1:0.7.

The molar ratio between the LNA monomer and the  $PN_2$ -reagent is typically in the range of 1:0.9 to 1:10, preferably 1:0.95 to 1:5, more preferably 1:1.




The currently most interesting LNA monomers are those which have a nucleobase selected from guanine, thymine, cytosine, methyl cytosine, uracil and adenine, typically in the protected form such as N-benzoyl protected cytosine ( $C^{Bz}$ ), N-benzoyl protected methyl cytosine ( $^{Me}C^{Bz}$ ), N-iso-butanoyl protected guanine ( $G^{Ibu}$ ), uracil (U), thymine (T), N-benzoyl protected adenine ( $A^{Bz}$ ).

## EXAMPLES

### EXAMPLE 1 - Phosphitylation methods

Four phosphitylation methods was examined and compared:



10                      Phosphitylation reaction. B =  $A^{Bz}$ ,  $C^{Bz}$ ,  $^{Me}C^{Bz}$ ,  $G^{Ibu}$  and T

- i.        2-Cyanoethyl-*N,N*-diisopropylchlorophosphoramidite, diisopropylethylamine
- ii.      2-Cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite, 1*H*-tetrazole
- iii.     2-Cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite, pyridinium trifluoroacetate
- 15 iv.     2-Cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite, 4,5-dicyanoimidazole

2-Cyanoethyl-*N,N*-diisopropylchlorophosphoramidite appeared to have several disadvantages compared to 2-cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite ( $PN_2$ -reagent). It is unstable at room temperature, expensive and difficult to handle due to its high reactivity. Furthermore undesired reactions with the nucleobases are often observed with 2-cyanoethyl-*N,N*-diisopropylchlorophosphoramidite (Figure 1).

Three different activators (1*H*-tetrazole, pyridinium trifluoroacetate and 4,5-dicyanoimidazole) were investigated with 2-cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite. 4,5-Dicyanoimidazole (DCI) gave the best results.

**EXAMPLE 2 - General experimental procedure for the preparation of LNA phosphoramidites 1-6:**

A 0.2 M solution of the LNA nucleoside (1.0 equiv) in anhyd  $\text{CH}_2\text{Cl}_2$  was stirred under argon. A 1.0 M solution of DCI (0.70 equiv) in anhyd MeCN was added followed by drop wise addition of 2-cyanoethyl-*N,N,N',N'*-tetraisopropylphosphorodiamidite (1.0 equiv). When the reaction was completed (monitored by analytical TLC\*) it was diluted with  $\text{CH}_2\text{Cl}_2$  and washed twice with sat. aq  $\text{NaHCO}_3$  and once with brine. The organic phase was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated *in vacuo* to give a colourless foam. All phosphoramidites were isolated in very high yields ( $\geq 95\%$ ). No by-products could be detected by HPLC-MS, TLC or  $^{31}\text{P}$ -NMR (purity  $\approx 100\%$ ). If starting materials are dry and pure chromatography is not necessary.

\*All phosphorylation reactions were completed in less than 2 h except for the G phosphoramidite (**2**) synthesis taking up to 4 h to be completed.

**Synthesis of (1*R*,3*R*,4*R*,7*S*)-3-(4-*N*-Benzoyl-5-methylcytosin-1-yl)-7-(2-cyanoethoxy-(diisopropylamino)phosphinoxy)-1-(4,4'-dimethoxytrityloxymethyl)-2,5-dioxabicyclo[2.2.1]heptane (**3**)**

(1*R*,3*R*,4*R*,7*S*)-3-(4-*N*-Benzoyl-5-methylcytosin-1-yl)-1-(4,4'-dimethoxytrityloxymethyl)-7-hydroxy-2,5-dioxabicyclo[2.2.1]heptane (4.86 g, 7.2 mmol) was dissolved in anhyd  $\text{CH}_2\text{Cl}_2$  (40 mL) and 4,5-dicyanoimidazole in MeCN (1.0 M, 5 mL) was added. Stirred at ambient temperature under argon. 2-Cyanoethyl-*N,N,N',N'*-tetraisopropylphosphorodiamidite (2.4 mL, 7.2 mmol) was added drop wise to the reaction mixture. After 2 h the reaction was diluted with  $\text{CH}_2\text{Cl}_2$  (100 mL) and transferred to a separation funnel and extracted with sat. aq.  $\text{NaHCO}_3$  (2x150 mL) and brine (150 mL). The combined aqueous phases were extracted with  $\text{CH}_2\text{Cl}_2$  (100 mL). The organic phases were pooled and dried ( $\text{Na}_2\text{SO}_4$ ). After filtration the organic phase was evaporated *in vacuo* to give a slightly yellow foam (6.0 g, 95% yield). A small sample was removed for comparison with the chromatographed material. The product was purified by Dry Column Vacuum Chromatography ( $\varnothing$  10 cm; Silica pretreated with 5%  $\text{Et}_3\text{N}$  in *n*-heptane v/v; 0→70% EtOAc, *n*-heptane v/v). Selected fractions containing **3** were combined and evaporated *in vacuo* to give a colourless foam (5.3 g, 84% yield).

No difference in coupling efficiency could be detected on the oligonucleotide synthesiser in a coupling efficiency assay (*vide supra*) between the crude and chromatographed phosphoramidite.

The compounds **1**, **2**, **4**, **5** and **6** were prepared in a similar manner.

Analytical data for phosphoramidites **1-6**.

- (1*R*,3*R*,4*R*,7*S*)-7-(2-Cyanoethoxy(diisopropylamino)phosphinoxy)-1-(4,4'-dimethoxytrityloxymethyl)-3-(thymine-1-yl)-2,5-dioxabicyclo[2.2.1]heptane (**1**)  
5 <sup>31</sup>P-NMR (CDCl<sub>3</sub>, 121.49 MHz): δ 150.0 (s), 149.3 (s).  
MS (ES): *m/z* calcd for C<sub>41</sub>H<sub>49</sub>N<sub>4</sub>O<sub>9</sub>P [M+H]<sup>+</sup>: 773.3. Found: 773.1.  
RP HPLC: R<sub>T</sub> = 5.89 min, 6.19 min.
- 10 (1*R*,3*R*,4*R*,7*S*)-7-(2-Cyanoethoxy(diisopropylamino)phosphinoxy)-1-(4,4'-dimethoxytrityloxymethyl)-3-(2-*N*-isobutylguanine-1-yl)-2,5-dioxabicyclo[2.2.1]heptane (**2**)  
<sup>31</sup>P-NMR (CDCl<sub>3</sub>, 121.49 MHz): δ 150.2 (s), 149.2 (s).  
MS (ES): *m/z* calcd for C<sub>45</sub>H<sub>54</sub>N<sub>7</sub>O<sub>9</sub>P [M+H]<sup>+</sup>: 868.4. Found: 868.0.  
RP HPLC: R<sub>T</sub> = 5.52 min, 5.72 min.
- 15 (1*R*,3*R*,4*R*,7*S*)-3-(4-*N*-Benzoyl-5-methylcytosine-1-yl)-7-(2-cyanoethoxy(diisopropylamino)phosphinoxy)-1-(4,4'-dimethoxytrityloxymethyl)-2,5-dioxabicyclo[2.2.1]heptane (**3**)  
<sup>31</sup>P-NMR (CDCl<sub>3</sub>, 121.49 MHz): δ 150.5 (s), 150.5 (s).
- 20 MS (ES): *m/z* calcd for C<sub>48</sub>H<sub>54</sub>N<sub>5</sub>O<sub>9</sub>P [M+H]<sup>+</sup>: 876.4. Found: 876.2.  
RP HPLC: R<sub>T</sub> = 7.67 min, 8.13 min.  
(HPLC solvent gradient: 0.0-0.5 min 95% B, 0.5-2.0 min 95%→100% B, 2.0-7.0 min 100% B, 7.0-7.5 min 100%→95% B, 7.5-12.0 min 95% B)
- 25 (1*R*,3*R*,4*R*,7*S*)-3-(6-*N*-Benzoyladenine-9-yl)-7-(2-cyanoethoxy(diisopropylamino)phosphinoxy)-1-(4,4'-dimethoxytrityloxymethyl)-2,5-dioxabicyclo[2.2.1]heptane (**4**)  
<sup>31</sup>P-NMR (CDCl<sub>3</sub>, 121.49 MHz): δ 150.1 (s), 149.7 (s).  
MS (ES): *m/z* calcd for C<sub>48</sub>H<sub>52</sub>N<sub>7</sub>O<sub>8</sub>P [M+H]<sup>+</sup>: 886.3. Found: 886.0.  
RP HPLC: R<sub>T</sub> = 6.65 min, 6.82 min.
- 30 (1*R*,3*R*,4*R*,7*S*)-7-(2-Cyanoethoxy(diisopropylamino)phosphinoxy)-1-(4,4'-dimethoxytrityloxymethyl)-5-*N*-methyl-3-(thymine-1-yl)-2-oxa-5-azabicyclo[2.2.1]heptane (**5**)  
<sup>31</sup>P-NMR (CDCl<sub>3</sub>, 121.49 MHz): δ 149.8 (s), 149.6 (s).  
MS (ES): *m/z* calcd for C<sub>42</sub>H<sub>52</sub>N<sub>5</sub>O<sub>8</sub>P [M+H]<sup>+</sup>: 786.3. Found: 786.2.
- 35 RP HPLC: R<sub>T</sub> = 5.87 min, 6.26 min.
- (1*S*,3*R*,4*S*,7*R*)-7-(2-Cyanoethoxy(diisopropylamino)phosphinoxy)-1-(4,4'-dimethoxytrityloxymethyl)-3-(thymine-1-yl)-2,5-dioxabicyclo[2.2.1]heptane (**6**)

$^{31}\text{P}$ -NMR ( $\text{CDCl}_3$ , 121.49 MHz):  $\delta$  150.9 (s), 150.6 (s).

MS (ES):  $m/z$  calcd for  $\text{C}_{41}\text{H}_{49}\text{N}_4\text{O}_9\text{P}$   $[\text{M}+\text{H}]^+$ : 773.3. Found: 773.1.

RP HPLC:  $R_T$  = 6.06 min, 6.44 min.

**CLAIMS**

1. A method for the synthesis of an LNA phosphoramidite, the method comprising phosphitylation of the 3'-OH groups of an LNA monomer with a 2-cyanoethyl-*N,N,N',N'*-tetra-substituted phosphoramidite in the presence of a nucleophilic activator.
- 5 2. The method according to claim 1, which comprises at the most one precipitation step.
3. The method according to claim 1 or 2, wherein the 2-cyanoethyl-*N,N,N',N'*-tetra-substituted phosphoramidite is selected from 2-cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite and 2-cyanoethyl-*N,N,N',N'*-tetraethylphosphoramidite.
- 10 4. The method according to any of the claims 1-3, wherein the nucleophilic activator is selected from 4,5-dicyanoimidazole (DCI), 4-alkylthioimidazole, 2-alkylthioimidazole, 2-nitroimidazole, 4-nitroimidazole, 4,5-dihaloimidazole, 4-haloimidazole, 2-haloimidazole and 5-alkoxytetrazole.
5. The method according to claims 1 or 2, wherein the 2-cyanoethyl-*N,N,N',N'*-tetra-  
15 substituted phosphoramidite is 2-cyanoethyl-*N,N,N',N'*-tetraisopropylphosphoramidite and the nucleophilic activator is 4,5-dicyanoimidazole.
6. The method according to any of the preceding claims, wherein the molar ratio between the LNA monomer and the nucleophilic activator is in the range of 1:0.0001 to 1:10.
- 20 7. The method according to any of the preceding claims, wherein the molar ratio between LNA monomer and the 2-cyanoethyl-*N,N,N',N'*-tetra-substituted phosphoramidite  $\text{PN}_2$ -reagent is in the range of 1:0.9 to 1:10.
8. The method according to any of the preceding claims, wherein the phosphitylation is allowed to proceed for 0.2-8 hours.
- 25 9. The method according to any of the preceding claims, wherein the LNA monomer has a nucleobase selected from guanine, thymine, cytosine, methyl cytosine, uracil and adenine.
10. The method according to any of the preceding claims, wherein the LNA monomer structure is selected from oxy- $\beta$ -D-ribo-LNA, thio- $\beta$ -D-ribo-LNA, amino- $\beta$ -D-ribo-LNA, oxy-  
30  $\alpha$ -L-ribo-LNA, thio- $\alpha$ -L-ribo-LNA and amino- $\alpha$ -L-ribo-LNA.

1/3








Figure 1



2/3

**1****2****3****4**

5

**Figure 2**

3/3



5

Figure 3



(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
23 January 2003 (23.01.2003)

PCT

(10) International Publication Number  
**WO 2003/006475 A3**

(51) International Patent Classification<sup>7</sup>: C07H 1/00, 19/02

(21) International Application Number:  
PCT/DK2002/000488

(22) International Filing Date: 12 July 2002 (12.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
PA 2001 01095 12 July 2001 (12.07.2001) DK

(71) Applicant (*for all designated States except US*): **Santaris Pharma A/S** [DK/DK]; Bøge Allé 3, DK-2970 Hørsholm (DK).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **KOCH, Troels** [DK/DK]; Funkiavej 47, DK-2300 Copenhagen S (DK). **ROSENBOHM, Christoph** [DK/DK]; Frederikssundsvej 55, 4. th., DK-2400 Copenhagen NV (DK). **PEDERSEN, Daniel, Sejer** [DK/DK]; Knabrostræde 22, 3., DK-1210 Copenhagen K (DK).

(74) Agent: **PLOUGMANN & VINGTOFT A/S**; Sundkrogs-  
gade 9, P.O. Box 831, DK-2100 Copenhagen Ø (DK).

(81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:  
26 February 2004

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: METHOD FOR PREPARATION OF LNA PHOSPHORAMIDITES

(57) Abstract: The present invention relates to large scale preparation of LNA phosphoramidites using a 2-cyanoethyl-N,N,N',N'-tetra-substituted phosphoramidite and a nucleophilic activator, e.g. 2-cyanoethyl-N,N,N',N'-tetraisopropylphosphoramidite and 4,5-dicyanoimidazole. The method is faster and more cost efficient than previously known methods.



WO 2003/006475 A3

# INTERNATIONAL SEARCH REPORT

In: tional Application No  
PCT/DK 02/00488

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07H1/00 C07H19/02

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SATOSHI OBIKA ET AL: "2'-0,4'-C-methylene bridged nucleic acid (2',4'-BNA): Synthesis and triplex-forming properties I" BIOORGANIC & MEDICINAL CHEMISTRY, vol. 9, 2001, pages 1001-1011, XP002902764 scheme 1, step j	1-10
X	SATOSHI OBIKA ET AL: "A 2', 4'-Bridged Nucleic Acid containing 2-Pyridone as a Nucleobase: Efficient Recognition of a C-G Interruption by Triplex Formation with a Pyrimidine Motif" ANGEW. CHEM. INT. ED., vol. 40, no. 11, 2001, pages 2079-2081, XP002902765 scheme 2, step e	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance  
"E" earlier document but published on or after the international filing date  
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
"O" document referring to an oral disclosure, use, exhibition or other means  
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.  
"&" document member of the same patent family

Date of the actual completion of the international search

22 October 2002

Date of mailing of the international search report

13. 02. 2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Viveca Norén

## INTERNATIONAL SEARCH REPORT

In International Application No  
PCT/DK 02/00488

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 16540 A (NEXSTAR PHARMACEUTICALS INC) 23 April 1998 (1998-04-23) scheme 3	1-10
X	--- SHOJI HAMAMOTO ET AL: "New Approach to the Synthesis of Deoxyribonucleoside Phosphoramidite Derivatives" CHEMISTRY LETTERS, vol. 8, 1986, pages 1401-1404, XP002902766 scheme 1	1-10
A	--- HAKANSSON A E ET AL: "CONVENIENT SYNTHESSES OF 7-HYDROXY-1-(HYDROXYMETHYL)-3-(THYMIN-1-YL)-2,5-DIOXABICYCLO2.2.1HEPTANES: ALPHA-6-RIBO- AND ALPHA-L-XYLO-CONFIGURED LNA NUCLEOSIDES" JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY. EASTON, US, vol. 65, no. 17, 25 August 2000 (2000-08-25), pages 5161-5166, XP000973191 ISSN: 0022-3263 scheme 2, step vi -----	1-10

## INTERNATIONAL SEARCH REPORT

### Information on patent family members

**Int**

**Additional Application No**

PCT/DK 02/00488

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 9816540	A	23-04-1998	AU	4800197 A		11-05-1998
			WO	9816540 A1		23-04-1998
-----						